편두통 유사 증상을 보인 갑상샘기능항진증 1례

한림대학교의료원 강남성심병원 소아청소년과

양원준 · 전상훈 · 오연정 · 이건희

A Case of Hyperthyroidism Mimicking Pediatric Migraine

Migraine is a common neurological disorder that often presents severe headache pain with nausea, vomiting, or sensitivity to light or sound. Children and adolescents with migraine experience drastically impaired quality of life. Therefore early diagnosis and treatments of headache are necessary in addition to accurate differentiation of secondary causes of headache. There are some reports that thyroid dysfunction especially hyperthyroidism may be responsible for producing headache symptoms. We detail a case of 15-year-old boy who was admitted to our hospital due to recurrent episodes of severe headache accompanied by nausea, compatible with the criteria for migraine without aura in the ICHD 3-beta, but finally diagnosed as hyperthyroidism. We further discuss the implications of current research regarding the association between thyroid dysfunction and headache. It is necessary to check the thyroid function tests in the pediatric patients experiencing unusual headache symptoms.

Key Words: Hyperthyroidism, Headache, Pediatric migraine, Thyroid dysfunction

Won Jun Yang, MD, Sang Hoon Chun, MD, Yeon Joung Oh, MD, Kon-Hee Lee, MD

Department of Pediatrics, Kangnam Sacred Heart Hospital, Hallym University College of Medicine, Seoul, Korea

Submitted: 25 August, 2016

Revised: 9 September, 2016

Accepted: 10 September, 2016

Correspondence to Kon-Hee Lee, MD
Department of Pediatrics, Kangnam Sacred Heart Hospital, Hallym University College of Medicine, 1 Singil-ro, Yeongdeungpo-gu, Seoul 07441, Korea
Tel: +82-2-829-5142, Fax: +82-2-829-5358
E-mail: headaches77@gmail.com

Introduction

Migraine is regarded as a common disorder in children and adolescents, with a prevalence of approximately 8.7%, according to a nationwide epidemiological study involving elementary, middle, and high school students in Korea¹⁾. Furthermore, a previous study involving nearly 1,200 children revealed that experiences of recurrent pain significantly impaired quality of life (QoL) during childhood²⁾. Therefore, early and accurate diagnoses as well as appropriate managements are critical in improving QoL in pediatric patients with migraine.

Diagnoses of migraine may be difficult in the patient with secondary causes of the headache attributed to uncommonly occurred disorders. In children and adolescents, the symptoms and signs of hyperthyroidism are similar to those in adults; fatigue, restlessness, tremors, enlarged thyroid, and exophthalmos occurs in up to one-third of pediatric patients; however, it is typically mild compared to adults³⁾.

Even though hypothyroidism was known to the causes of secondary headache, but there a few reports that hyperthyroidism may cause headache or aggravates previously existing migraine^{4,5)}.

In the present report, we detail the case of an adolescent patient exhibiting symptoms of migraine due to hyperthyroidism.

Case report

A 15-year-old boy was admitted to our pediatric department for evaluation of severe, frequent headaches. He had been healthy 3 weeks before admission. Headaches occurred mainly in the frontal or temporal parts of the head, approximately four times per week, with each episode lasting about 2–3 hours. The patient described the pain as ‘throbbing’ and rated the intensity of pain as 7–8 on a visual analog scale. Headache pain became aggravated with normal walking, and was associated with nausea, but no symptoms of photophobia or phonophobia. During headache attacks, the patient experienced transient dizziness or blurred vision lasting for about 30 seconds six to seven times.

On the physical examination, the patient displayed no evidence of hyperthyroidism such as tachycardia, protruded eyeballs or enlarged thyroid gland. On the ophthalmoscopic examination (PanOptic™ Ophthalmoscope, Welch Allyn), there were no findings of papilledema or widening of optic cup etc. Other neurologic examinations including mental status, functions of cranial nerve, motor and sensory nerve systems were normal.

The patient reported no history of head trauma or other health problems, recently unusual weight loss, and drug overuse or smoking. The patient’s mother was diagnosed with diabetes and reported occasional experiences of headaches with nausea.

The patient’s blood pressure and heart rate at the time of evaluation were 110/70 mmHg and 83 beats per minute, respectively. Blood test revealed white blood cell count of 5,740/ μ L; platelet count of 265,000/ μ L; hemoglobin level of 15.3 g/dL; and normal hematocrit of 44.4%. Additional analyses showed no

abnormal findings (AST/ALT=26/26 IU/L; BUN/Cr=2.9/0.75 mg/dL; glucose=88 mg/dL; Na=139 mMol/L; K=4.5 mMol/L; and Cl = 101 mMol/L). Fluorescent antinuclear antibody (FANA) testing was negative.

However, hormone analysis yielded decreased levels of thyroid-stimulating hormone (TSH) (0.076 μ IU/mL; normal range=0.38–4.7 μ IU/mL), as well as increased levels of T3 (>800 ng/dL; normal range=59–174 ng/dL) and free T4 (2.19 ng/dL; normal range=0.89–1.76 ng/dL). Anti-thyroglobulin levels were also increased (202.5 IU/mL; normal range=5–100 IU/mL). Thyroid-stimulating antibody (TSAb) and antithyroid microsomal antibody results were negative.

On the ultrasound examination of the neck, abnormal findings of decreased and heterogeneous echogenicity were appeared on the thyroid (Fig. 1). Thyroid images were obtained using a pinhole gamma camera 10 min after injection of Tc-99m pertechnetate 370 MBq (10 mCi). Isotope uptake in the bilateral thyroid had decreased to 1.2%, with relative increases observed in the surrounding soft tissues (Fig. 2).

Brain MRI with diffusion weighted imaging revealed unremarkable findings. Clinical psychological screening tests such as Spielberger’s State- and Trait Anxiety Inventory for Children, Kovacs Children’s Depression Inventory and Korean Child Behavior Check List were performed, and showed no abnormal findings such as anxiety, depression or behavioral problems. Cerebrospinal fluid (CSF) test was not performed because of refusal by the patient’s parent.

Initially the patient was diagnosed with ‘migraine without aura’, which was compatible to international criteria of headache

Fig. 1. Sonographic examination revealed decreased and heterogeneous echogenicity in both thyroid lobes.

Fig. 2. Thyroid scan showed isotope uptake in the bilateral thyroid had decreased to 1.2%, with relative increases observed in the surrounding soft tissues.

disorders III beta (ICHD 3-beta), but finally as 'secondary headache due to hyperthyroidism'. He was discharged with initiation of daily treatment with anti-thyroid drug (methimazole, 15 mg daily). After 2 weeks of drug therapy, serum levels of thyroid hormone returned to normal level, and his headache symptoms had improved. The patient had experienced no recurrence of headache during the follow-up period of 6 months.

Discussion

It is still controversial whether headache attributed to thyroid dysfunction or not⁵⁻⁸. ICHD 3-beta describes '10.4 headache attributed to hypothyroidism' on the secondary causes of the headache attributed to disorder of homeostasis⁴. In a study of 21 chronic headache patients, none of hypothyroidism was occurred⁹, but bilaterally continuous, non-pulsatile headaches were found about 30% of 102 patients with hypothyroidism¹⁰.

The mechanisms between hypothyroidism and headaches are unknown, but it is suggested that low levels of thyroid hormones may induce pro-nociceptive state⁵. On the purinergic system, adenosine has anti-nociceptive properties, while adenosine triphosphate (ATP) has pro-nociceptive actions in different pain models. In the study of rats, ATP hydrolysis was decreased in the spinal cord of hypothyroid rats in parallel with a significant increase in nociceptive response. Therefore, hypothyroidism may involve adenine nucleotides of purinergic system and significantly influence on the modulation of pain pathways¹¹.

On the other hand, hyperthyroidism is mentioned under the appendix heading 'A10.8.2 headache attributed to other metabolic

or systemic disorder⁴. Three cases with thyrotoxicosis presented with severe headache with unremitting quality, improved following anti-thyroid drugs¹². Unexplained headache with vomiting and abnormal liver function for 3 weeks was diagnosed as hyperthyroidism after re-admission¹³. Hyperthyroidism caused by tumors or autonomic dysreflexia may present severe headache^{14,16}.

Pathophysiology of headaches with hyperthyroidism is poorly understood. Recently proposed mechanisms that thyroid hormone may affect anti-diuretic hormone related free water clearance, would lead to increase pressure¹⁷. Some reports that hyperthyroidism with headache showed findings of increased intracranial pressure (ICP)^{18,19}. Another mechanism was that hyperthyroidism increased coagulation factors²⁰ and caused cerebral venous thrombosis, which elevated ICP as mechanical factors^{17,21}.

In our case, CSF study was not done, but there were no signs of increased ICP with signs of papilledema in the ophthalmoscopic test and findings of dilated ventricle, protrusion of the optic disc or widening of the optic nerve sheath in MRI²². Our patient showed symptoms of transient visual blurring, which was different aspects of occipital seizures, producing a sudden onset of positive visual loss with multiple, brightly colored, small circular spots, circles, or balls²³. Electroencephalography (EEG) generally does not give us additional information in migraine patients, but is considered for excluding disorders such as migraine-triggered seizure, or recurrent episodes of confusion²⁴.

In the examination of headache patients, it still questionable that routine test of thyroid function is needed because of thyroid dysfunction is not common among the pediatric migraine^{25,26}. But tests of thyroid function are necessary in the unusual cases of headache or new onset of headache without previous headache history^{27,28}.

요약

편두통은 소아청소년에서 흔한 질환으로 조기에 정확한 진단과 적절한 치료가 환자의 삶의 질을 높이는데 매우 중요하다. 정확한 진단을 위해서 두통의 2차적 원인을 감별해야 하는데, 매우 드문 질환이 원인인 경우는 어려움이 따른다. 최근 2차적 원인 중에서 갑상샘기능이상인 드물게 두통과 관련이 있다는 보고가 있다. 저자들은 갑상샘기능항진증으로 인하여 편두통 증상을 보인 환자 증례를 치험하였기에 보고한다. 15세 남아가 2주전부터 처음으로 시작된 주당 4회 정도의 심한 두통(7-8시간 아날로그 척도)을 호소하며 외래 통해 입원하였다. 두통의 통증 부위는 주로 앞이마 부위로 2-3시간 정도 지속되었으며 구역 증세도 동반하였다. 두통 동안에 30초 정도의 어지러움과 시야흐림 증상이 있었다. 생체 징후는 안정적이었으며 뇌자기공명영상 검사에서 특이소견 보이지 않았다. 말초혈액 검

사에서 특이소견 보이지 않았으나 호르몬 검사에서 갑상샘 자극호르몬 수치가 0.076 μ U/mL로 증가되어 있었고, T3 >800 ng/dL, free T4 2.19 ng/dL로 증가되어 있었다. 갑상샘 초음파 검사에서 불균일한 저에코양상을 보였으며, 핵의학 검사에서는 양측 갑상샘 동위 원소 섭취율이 1.2%로 감소되어 있었다. 최초로 임상적으로 무조짐편 두통을 의심하였으나, 갑상샘기능 항진증 진단 하에 methimazole (15 mg/day)로 치료를 시작하였다. 15일간 투약 후 두통증세 호전되었으며 갑상샘 호르몬 기능도 정상으로 돌아왔다. 소아두통환자에서 설명되는 않는 두통 설명되지 않는 두통 특히 만성 혹은 두통 병력이 없는 환자가 갑자기 심한 두통을 호소하는 경우 등의 환자에서는 갑상샘 기능검사에 대한 검토도 필요할 것으로 생각된다.

References

- 1) Rho YI, Chung HJ, Lee KH, Eun BL, Eun SH, Nam SO, et al. Prevalence and clinical characteristics of primary headaches among school children in South Korea: a nationwide survey. *Headache* 2012;52:592-9.
- 2) Wöber-Bingöl Ç, Wöber C, Uluduz D, Uygunoğlu U, Aslan TS, Kernmayer M, et al. The global burden of headache in children and adolescents-developing a questionnaire and methodology for a global study. *J Headache Pain* 2014;15:86.
- 3) Hanley P, Lord K, Bauer AJ. Thyroid Disorders in Children and Adolescents: A Review. *JAMA Pediatr* 2016.
- 4) Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders, 3rd ed. beta version. *Cephalalgia* 2013;33:629-808.
- 5) Tepper DE, Tepper SJ, Sheftell FD, Bigal ME. Headache attributed to hypothyroidism. *Curr Pain Headache* 2007;11:304-9.
- 6) Larner AJ. Thyroid dysfunction and headache. *J Headache Pain* 2006;7:51-2.
- 7) Gipponi S1, Venturelli E, Rao R, Liberini P, Padovani A. Hypertension is a factor associated with chronic daily headache. *Neurol Sci* 2010;31 Suppl 1:S171-3.
- 8) Lima Carvalho MF, de Medeiros JS, Valença MM. Headache in recent onset hypothyroidism: Prevalence, characteristics and outcome after treatment with levothyroxine. *Cephalalgia* 2016; 43:1-7.
- 9) Amy JR. Tests of thyroid function in chronic headache patients. *Headache* 1987;27:351-3.
- 10) Moreau T, Manceau E, Giroud-Baleyrier F, Dumas R, Giroud M. Headache in hypothyroidism. Prevalence and outcome under thyroid hormone therapy. *Cephalalgia* 1998;18:687-9.
- 11) Bruno AN, Pochmann D, Ricachenevsky FK, Fontella FU, Bonan CD, Dalmaz C, et al. Nociceptive response and adenine nucleotide hydrolysis in synaptosomes isolated from spinal cord of hypothyroid rats. *Neurochem Res* 2005;30:1155-61.
- 12) Stone J, Foulkes A, Adamson K, Stevenson L, Al-Shahi Salman R. Thyrotoxicosis presenting with headache. *Cephalalgia* 2007;27: 561-2.
- 13) Ashawesh K, Abdulqawi R, Ahmad S. Hyperthyroidism presenting with persistent vomiting, headache and deranged liver function tests. *Eur J Intern Med* 2008;19:72.
- 14) Arnason T, Clarke DB, Imran SA. Hyperthyroidism caused by a pituitary adenoma. *CMAJ* 2011;183:E757.
- 15) Olt S, Şirik M. TSH secreting adenoma: a rare cause of severe headache. *Pan Afr Med J* 2016;23:2.
- 16) Chang JH, Han JE, Sohn HJ, Shin YB, Ko HY. Hyperthyroidism as a cause of autonomic dysreflexia. *Am J Phys Med Rehabil* 2008; 87:768-72.
- 17) Sheldon CA, Kwon YJ, Liu GT, McCormack SE. An integrated mechanism of pediatric pseudotumor cerebri syndrome: evidence of bioenergetic and hormonal regulation of cerebrospinal fluid dynamics. *Pediatr Res* 2015;77:282-9.
- 18) Herwig U, Sturzenegger M. Hyperthyroidism mimicking increased intracranial pressure. *Headache* 1999;39:228-30.
- 19) Yaka E, Cakmur R. Increased intracranial pressure due to hyperthyroidism. *Cephalalgia* 2010;30:878-80.
- 20) Poole-Wilson PA, Langer GA. Effect of pH on ionic exchange and function in rat and rabbit myocardium. *Am J Physiol* 1975;229: 570-81.
- 21) Pekdemir M, Yilmaz S, Ersel M, Sarisoy HT. A rare cause of headache: cerebral venous sinus thrombosis due to hyperthyroidism. *Am J Emerg Med* 2008;26:383.e1-2.
- 22) Passi N1, Degnan AJ, Levy LM. MR imaging of papilledema and visual pathways: effects of increased intracranial pressure and pathophysiologic mechanisms. *AJNR Am J Neuroradiol* 2013;34: 919-24.
- 23) Thurtel MJ, Rucker JC. Transient visual loss. *Int Ophthalmol Clin* 2009;49:147-66.
- 24) Srinivasa R, Kumar R. Migraine variants and beyond. *J Assoc Physicians India* 2010;58 Suppl:14-7.
- 25) Dilek Toprak, Kemal Demirkıran, Hülya Ellidokuz. Is it important to test thyroid function tests in migraineurs?. *TJFMPC* 2007; 4:47-51.
- 26) Ekici B, Cebeci AN. The debate on the link between subclinical hypothyroidism and childhood migraine: is initial endocrinological evaluation necessary for children with migraine? *Acta Neurol Belg* 2015;115:123-7.
- 27) De Luca GC, Bartleson JD. When and how to investigate the patient with headache. *Semin Neurol* 2010;30:131-44.
- 28) Kim HS, Shin SH, Hwang IT, Lee KH. A case of chronic daily headache attributed to Hashimoto's thyroiditis. *J Korean Child Neurol Soc* 2007;15:111-5.